


In the Claims:

The current status of all claims is listed below and supercedes all previous lists of claims.

Please amend claims 1-20, and add new claims 21-28 as follows.

1. (currently amended) A compound of formula (I):



(I)

wherein:

R^1 is hydrogen, C_{1-6} alkyl, C_{3-6} cycloalkyl or aryl; wherein said C_{1-6} alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, carbamoyl, carboxy, C_{1-6} alkoxy, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl) $_2$ amino, ~~C_{1-6} alkylcarbamoyl~~ C_{1-6} alkylcarbamoyl, C_{1-6} alkylS(O) $_a$ wherein a is 0-2, C_{3-6} cycloalkyl or aryl; and wherein any aryl group may be optionally substituted by one or two substituents selected from halo, hydroxy, C_{1-6} alkyl or C_{1-6} alkoxy;

R^2 and R^5 are independently hydrogen, a branched or unbranched C_{1-6} alkyl, C_{3-6} cycloalkyl or aryl; wherein said C_{1-6} alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, cyano, carbamoyl, carboxy, C_{1-6} alkoxy, aryl C_{1-6} alkoxy, $(C_1-C_4)_3Si$, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl) $_2$ amino, ~~C_{1-6} alkylS(O)~~, C_{1-6} alkylS(O) $_a$, C_{3-6} cycloalkyl, aryl or aryl C_{1-6} alkylS(O) $_a$, wherein a is 0-2; and wherein any aryl group may be optionally substituted by one or two substituents selected from halo, hydroxy, C_{1-6} alkyl or C_{1-6} alkoxy;

R^3 is hydrogen, alkyl, halo, C_{1-6} alkoxy or C_{1-6} alkylS-;

R^4 is hydrogen, C_{1-6} alkyl, halo or C_{1-6} alkoxy;


R^6 is hydrogen, C_{1-6} alkyl, or aryl C_{1-6} alkyl;

wherein R⁵ and R² may form a ring with 2-7 carbon atoms and wherein R⁶ and R² may form a ring with 3-6 carbon atoms;

or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof;

with the proviso that said compound is not 3-(R)-4-(R)-1-(phenyl)-3-[2-(4-fluorophenyl)-2-hydroxyethylsulphanyl]-4-[4-(N-{N-[(R)-1-(carboxy)-2-(hydroxy)ethyl]carbamoylmethyl}carbamoylmethoxy)phenyl]azetidin-2-one; or 3-(R)-4-(R)-1-(phenyl)-3-[2-(4-fluorophenyl)-2-hydroxyethylsulphanyl]-4-{4-[N-((R)- α -{N-[(S)-1-(carboxy)-2-(hydroxy)ethyl]carbamoyl}benzyl)carbamoylmethoxy]phenyl}azetidin-2-one.

2. (currently amended) A compound of formula (I2):



(I2)

wherein:

R¹ is hydrogen, C₁₋₆alkyl, C₃₋₆cycloalkyl or aryl; wherein said C₁₋₆alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, carbamoyl, carboxy, C₁₋₆alkoxy, N-(C₁₋₆alkyl)amino, N,N-(C₁₋₆alkyl)₂amino, ~~C₁₋₆alkylcarbonylamino~~ C₁₋₆alkylcarbonylamino, C₁₋₆alkylS(O)_a wherein a is 0-2, C₃₋₆cycloalkyl or aryl; and wherein any aryl group may be optionally substituted by one or two substituents selected from halo, hydroxy, C₁₋₆alkyl or C₁₋₆alkoxy;

R² and R⁵ are independently hydrogen, a branched or unbranched C₁₋₆alkyl, C₃₋₆cycloalkyl or aryl; wherein said C₁₋₆alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, cyano, carbamoyl, carboxy, C₁₋₆alkoxy, aryl C₁₋₆alkoxy, (C₁₋₄)₃Si, N-(C₁₋₆alkyl)amino, N,N-(C₁₋₆alkyl)₂amino, ~~C₁₋₆alkylS(O)~~ C₁₋₆alkylS(O)_a, C₃₋₆cycloalkyl, aryl or

aryl C₁₋₆ alkylS(O)_a, wherein a is 0-2; and wherein any aryl group may be optionally substituted by one or two substituents selected from halo, hydroxy, C₁₋₆alkyl or C₁₋₆alkoxy;

R³ is hydrogen, alkyl, halo, C₁₋₆alkoxy or C₁₋₆ alkylS-;

R⁴ is hydrogen, C₁₋₆ alkyl, halo or C₁₋₆alkoxy;

R⁶ is hydrogen, C₁₋₆ alkyl, or arylC₁₋₆ alkyl;

wherein R⁵ and R² may form a ring with 2-7 carbon atoms and wherein R⁶ and R² may form a ring with 3-6 carbon atoms;

or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof;

with the proviso that said compound is not 3-(R)-4-(R)-1-(phenyl)-3-[2-(4-fluorophenyl)-2-hydroxyethylsulphanyl]-4-[4-(N-{N-[(R)-1-(carboxy)-2-(hydroxy)ethyl]carbamoylmethyl}carbamoylmethoxy)phenyl]azetidin-2-one; or 3-(R)-4-(R)-1-(phenyl)-3-[2-(4-fluorophenyl)-2-hydroxyethylsulphanyl]-4-{4-[N-((R)-α-{N-[(S)-1-(carboxy)-2-(hydroxy) ethyl]carbamoyl}benzyl)carbamoylmethoxy]phenyl}azetidin-2-one.

3. (currently amended) A compound according to claim 1 ~~or 2~~, wherein:

R¹ is hydrogen or phenyl.

4. (currently amended) A compound according to ~~any of the preceding claims~~ claim 1, wherein:

R² is hydrogen, a branched or unbranched C₁₋₆alkyl, C₃₋₆cycloalkyl or aryl; wherein said C₁₋₆alkyl may be optionally substituted by one or more hydroxy, amino, acylamino, C₁₋₆alkylS(O)_a wherein a is 0-2, C₃₋₆cycloalkyl or aryl; and wherein any aryl group may be optionally substituted by hydroxy, alkyl, alkoxy or cyano.

5. (currently amended) A compound according to ~~any of the preceding claims~~ claim 1, wherein:

R³ is hydrogen, C₁-C₂alkyl, halo or methoxy.

6. (currently amended) A compound according to ~~any of the preceding claims~~ claim 1, wherein:

R^3 is hydrogen, methyl, chlorine, fluorine, C_{1-6} alkylS-, or methoxy.

7. (currently amended) A compound according to ~~any of the preceding claims~~ claim 1, wherein:

R^4 is hydrogen or halo.

8. (currently amended) A compound according to ~~any of the preceding claims~~ claim 1, wherein:

R^4 is chlorine or fluorine.

9. (currently amended) A compound according to ~~any of the preceding claims~~ claim 1, wherein:

R^6 is hydrogen, C_{1-6} alkyl, aryl C_{1-6} alkyl or R^6 and R^2 form a ring with 3-6 carbon atoms.

10. (currently amended) A compound according to claim 1, wherein:

R^1 is hydrogen;

R^2 is a branched or unbranched C_{1-4} alkyl, optionally substituted by a C_{3-6} cycloalkyl, alkylS-, aryl optionally substituted by hydroxy or cyano, amino, *N*-(C_{1-6} alkyl)amino, *N,N*-(C_{1-6} alkyl)₂amino or aryl C_{1-6} alkylS(O)_a, wherein a is ~~0-2~~ 0-2;

R^3 and R^4 are halo;

R^5 is hydrogen or C_{1-6} alkyl; and

R^6 is hydrogen.

11. (currently amended) One or more compounds chosen from:

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-*N*⁶-acetyl-D-lysine;

1-(4-Fluorophenyl)-3-(*R*)-[2-(4-fluorophenyl)-2-hydroxyethylthio]-4-(*R*)-{4-[*N*-{*N*-[2-

(phenyl)-1-(R)-(carboxy)ethyl]carbamoylmethyl}carbamoylmethoxy]phenyl}azetidin-2-one;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-valine;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-tyrosine;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-proline;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-lysine;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-hydroxy-2-(4-methoxyphenyl)ethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-valine;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-2-butylnorleucine;

N-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-*S*-methyl-L-cysteine;

N-{[4-((2*R*,3*R*)-1-(4-chlorophenyl)-3-{[2-(4-chlorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-3-cyclohexyl-D-alanine;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-3-cyclohexyl-D-alanine;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-4-methylleucine;

N-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-L-alanyl-D-valine;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-hydroxy-2-(4-methylphenyl)ethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-valine;

N-{[4-((2*R*,3*R*)-1-(4-chlorophenyl)-3-{[2-(4-chlorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-valine;

N-{[4-((2*R*,3*R*)-1-(4-chlorophenyl)-3-{[2-(4-chlorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-3-methyl-D-valine;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-

oxoazetidin-2-yl)phenoxy]acetyl}glycyl-3-(2-naphthyl)-D-alanine;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-3-methyl-D-valine;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-(3*R*,4*S*,5*R*)-3,4,5,6-tetrahydroxy-D-~~norleucine~~.

norleucine;

N-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-*N*,2-~~dimethylalanine~~ dimethylalanine;


N-({4-[(2*R*,3*R*)-1-(4-Fluorophenyl)-3-({2-hydroxy-2-[4-(methylthio)phenyl]ethyl}thio)-4-oxoazetidin-2-yl]phenoxy}acetyl}glycyl-3-methyl-D-~~valine~~ valine;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-*S*-(4-methylbenzyl)-D-~~cysteine~~ cysteine;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-*S*-(*tert*-butyl)-D-~~cysteine~~ cysteine; and

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-b,b-dimethyl-D-phenylalanine.

12. (currently amended) A compound of the formula (XV) or hydrolysable esters or amides thereof:



(XV)

wherein:

R^1 is hydrogen, C_{1-6} alkyl, C_{3-6} cycloalkyl or aryl; wherein said C_{1-6} alkyl may be

optionally substituted by one or more hydroxy, amino, guanidino, carbamoyl, carboxy, C₁₋₆alkoxy, *N*-(C₁₋₆alkyl)amino, *N,N*-(C₁₋₆alkyl)₂amino, ~~C₁₋₆alkylcarboxylamino~~ C₁₋₆alkylcarboxylamino, C₁₋₆alkylS(O)_a wherein a is 0-2, C₃₋₆cycloalkyl or aryl; and wherein any aryl group may be optionally substituted by one or two substituents selected from halo, hydroxy, C₁₋₆alkyl or C₁₋₆alkoxy;

R² and R⁵ are independently hydrogen, a branched or unbranched C₁₋₆alkyl, C₃₋₆cycloalkyl or aryl; wherein said C₁₋₆alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, carbamoyl, carboxy, C₁₋₆alkoxy, aryl C₁₋₆alkoxy, (C₁₋₄)₃Si, *N*-(C₁₋₆alkyl)amino, *N,N*-(C₁₋₆alkyl)₂amino, C₁₋₆alkylS(O)_a, ~~aryl C₁₋₆alkylS(O)_a~~, aryl C₁₋₆alkylS(O)_{a2}, wherein a is 0-2, C₃₋₆cycloalkyl or aryl; and wherein any aryl group may be optionally substituted by one or two substituents selected from halo, hydroxy, C₁₋₆alkyl or C₁₋₆alkoxy;

R³ is hydrogen, alkyl, halo, C₁₋₆alkoxy or C₁₋₆ alkylS-;

R⁴ is hydrogen, C₁₋₆ alkyl, halo or C₁₋₆alkoxy;

R⁶ is hydrogen, C₁₋₆ alkyl, or arylC₁₋₆ alkyl; and

R⁷ is an hydroxy group or a C₁₋₃ alkoxy group;

wherein R⁵ and R² may form a ring with 2-7 carbon atoms and wherein R⁶ and R² may form a ring with 3-6 carbon atoms;

or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof;

with the proviso that said compound is not 3-(R)-4-(R)-1-(phenyl)-3-[2-(4-fluorophenyl)-2-hydroxyethylsulphanyl]-4-[4-(*N*-{*N*-[(R)-1-(carboxy)-2-(hydroxy)ethyl]carbamoylmethyl}carbamoylmethoxy)phenyl]azetidin-2-one; or 3-(R)-4-(R)-1-(phenyl)-3-[2-(4-fluorophenyl)-2-hydroxyethylsulphanyl]-4-{4-[*N*-((R)-α-{*N*-[(S)-1-(carboxy)-2-(hydroxy) ethyl]carbamoyl}benzyl)carbamoylmethoxy]phenyl}azetidin-2-one.

13. (currently amended) A method of treating or preventing a hyperlipidemic condition ~~hyperlipidemic conditions~~ comprising the administration of an effective amount of a compound according to ~~any one of claims 1 to 12~~ claim 1 to a mammal in need thereof.


14. (currently amended) A method of treating or preventing atherosclerosis comprising the administration of an effective amount of a compound according to ~~any one of claims 1 to 12~~ claim 1 to a mammal in need thereof.

15. (currently amended) A method for treating or preventing Alzheimers' disease comprising the administration of an effective amount of a compound according to ~~any one of claims 1 to 12~~ claim 1 to a mammal in need thereof.

16. (currently amended) A method for treating or preventing a cholesterol associated tumor ~~cholesterol associated tumors~~ comprising the administration of an effective amount of a compound according to ~~any one of claims 1 to 12~~ claim 1 to a mammal in need thereof.

17. (currently amended) A pharmaceutical formulation comprising a compound according to ~~any one of claims 1 to 12~~ claim 1 in admixture with a pharmaceutically acceptable adjuvant, diluent and/or carrier ~~adjuvants, diluents and/or carriers~~.

18. (currently amended) A combination of a compound according to formula (I)



(I)

wherein:

R¹ is hydrogen, C₁₋₆alkyl, C₃₋₆cycloalkyl or aryl; wherein said C₁₋₆alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, carbamoyl, carboxy, C₁₋₆alkoxy, N-(C₁₋₆alkyl)amino, N,N-(C₁₋₆alkyl)₂amino, C₁-C₆alkylcarbonylamino, C₁₋₆alkylS(O)_a wherein a is 0-2, C₃₋₆cycloalkyl or aryl; and wherein any aryl group may be

optionally substituted by one or two substituents selected from halo, hydroxy, C₁₋₆alkyl or C₁₋₆alkoxy;

R² and R⁵ are independently hydrogen, a branched or unbranched C₁₋₆alkyl, C₃₋₆cycloalkyl or aryl; wherein said C₁₋₆alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, cyano, carbamoyl, carboxy, C₁₋₆alkoxy, aryl C₁₋₆alkoxy, (C₁₋₄)₃Si, N-(C₁₋₆alkyl)amino, N,N-(C₁₋₆alkyl)₂amino, C₁₋₆alkylS(O)_a, C₃₋₆cycloalkyl, aryl or aryl C₁₋₆alkylS(O)_a, wherein a is 0-2; and wherein any aryl group may be optionally substituted by one or two substituents selected from halo, hydroxy, C₁₋₆alkyl or C₁₋₆alkoxy;


R³ is hydrogen, alkyl, halo, C₁₋₆alkoxy or C₁₋₆alkylS-;

R⁴ is hydrogen, C₁₋₆alkyl, halo or C₁₋₆alkoxy;

R⁶ is hydrogen, C₁₋₆alkyl, or arylC₁₋₆alkyl;

wherein R⁵ and R² may form a ring with 2-7 carbon atoms and wherein R⁶ and R² may form a ring with 3-6 carbon atoms;

or according to formula (I2)



(I2)

wherein:

R¹ is hydrogen, C₁₋₆alkyl, C₃₋₆cycloalkyl or aryl; wherein said C₁₋₆alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, carbamoyl, carboxy, C₁₋₆alkoxy, N-(C₁₋₆alkyl)amino, N,N-(C₁₋₆alkyl)₂amino, C₁₋₆alkylcarbonylamino, C₁₋₆alkylS(O)_a wherein a is 0-2, C₃₋₆cycloalkyl or aryl; and wherein any aryl group may be optionally substituted by one or two substituents selected from halo, hydroxy, C₁₋₆alkyl or C₁₋₆alkoxy;

R² and R⁵ are independently hydrogen, a branched or unbranched C₁₋₆alkyl,

C₃₋₆cycloalkyl or aryl; wherein said C₁₋₆alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, cyano, carbamoyl, carboxy, C₁₋₆alkoxy, aryl C₁₋₆alkoxy, (C₁₋₄)₃Si, N-(C₁₋₆alkyl)amino, N,N-(C₁₋₆alkyl)₂amino, C₁₋₆alkylS(O)_a, C₃₋₆cycloalkyl, aryl or aryl C₁₋₆alkylS(O)_a, wherein a is 0-2; and wherein any aryl group may be optionally substituted by one or two substituents selected from halo, hydroxy, C₁₋₆alkyl or C₁₋₆alkoxy;

R³ is hydrogen, alkyl, halo, C₁₋₆alkoxy or C₁₋₆alkylS-;


R⁴ is hydrogen, C₁₋₆alkyl, halo or C₁₋₆alkoxy;

R⁶ is hydrogen, C₁₋₆alkyl, or arylC₁₋₆alkyl;

wherein R⁵ and R² may form a ring with 2-7 carbon atoms and wherein R⁶ and R² may form a ring with 3-6 carbon atoms;

with a PPAR alpha and/or gamma agonist.

19. (currently amended) A combination of a compound according to formula (I)



(I)

wherein:

R¹ is hydrogen, C₁₋₆alkyl, C₃₋₆cycloalkyl or aryl; wherein said C₁₋₆alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, carbamoyl, carboxy, C₁₋₆alkoxy, N-(C₁₋₆alkyl)amino, N,N-(C₁₋₆alkyl)₂amino, C₁₋₆alkylcarbonylamino, C₁₋₆alkylS(O)_a wherein a is 0-2, C₃₋₆cycloalkyl or aryl; and wherein any aryl group may be optionally substituted by one or two substituents selected from halo, hydroxy, C₁₋₆alkyl or C₁₋₆alkoxy;

R² and R⁵ are independently hydrogen, a branched or unbranched C₁₋₆alkyl, C₃₋₆cycloalkyl or aryl; wherein said C₁₋₆alkyl may be optionally substituted by one or more

hydroxy, amino, guanidino, cyano, carbamoyl, carboxy, C₁₋₆alkoxy, aryl C₁₋₆alkoxy, (C₁₋₄)₃Si, N-(C₁₋₆alkyl)amino, N,N-(C₁₋₆alkyl)₂amino, C₁₋₆alkylS(O)_a, C₃₋₆cycloalkyl, aryl or aryl C₁₋₆alkylS(O)_a, wherein a is 0-2; and wherein any aryl group may be optionally substituted by one or two substituents selected from halo, hydroxy, C₁₋₆alkyl or C₁₋₆alkoxy;


R³ is hydrogen, alkyl, halo, C₁₋₆alkoxy or C₁₋₆alkylS-;

R⁴ is hydrogen, C₁₋₆alkyl, halo or C₁₋₆alkoxy;

R⁶ is hydrogen, C₁₋₆alkyl, or arylC₁₋₆alkyl;

wherein R⁵ and R² may form a ring with 2-7 carbon atoms and wherein R⁶ and R² may form a ring with 3-6 carbon atoms;

or according to formula (I2)



(I2)

wherein:

R¹ is hydrogen, C₁₋₆alkyl, C₃₋₆cycloalkyl or aryl; wherein said C₁₋₆alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, carbamoyl, carboxy, C₁₋₆alkoxy, N-(C₁₋₆alkyl)amino, N,N-(C₁₋₆alkyl)₂amino, C₁₋₆alkylcarbonylamino, C₁₋₆alkylS(O)_a wherein a is 0-2, C₃₋₆cycloalkyl or aryl; and wherein any aryl group may be optionally substituted by one or two substituents selected from halo, hydroxy, C₁₋₆alkyl or C₁₋₆alkoxy;

R² and R⁵ are independently hydrogen, a branched or unbranched C₁₋₆alkyl, C₃₋₆cycloalkyl or aryl; wherein said C₁₋₆alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, cyano, carbamoyl, carboxy, C₁₋₆alkoxy, aryl C₁₋₆alkoxy, (C₁₋₄)₃Si, N-(C₁₋₆alkyl)amino, N,N-(C₁₋₆alkyl)₂amino, C₁₋₆alkylS(O)_a, C₃₋₆cycloalkyl, aryl or aryl C₁₋₆alkylS(O)_a, wherein a is 0-2; and wherein any aryl group may be optionally substituted by one or

two substituents selected from halo, hydroxy, C₁₋₆alkyl or C₁₋₆alkoxy;

R³ is hydrogen, alkyl, halo, C₁₋₆alkoxy or C₁₋₆alkylS-;

R⁴ is hydrogen, C₁₋₆alkyl, halo or C₁₋₆alkoxy;


R⁶ is hydrogen, C₁₋₆alkyl, or arylC₁₋₆alkyl;

wherein R⁵ and R² may form a ring with 2-7 carbon atoms and wherein R⁶ and R² may form a ring with 3-6 carbon atoms;

with an HMG Co-A reductase inhibitor.


20. (currently amended) A process for preparing a compound of formula (I) or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof ~~which process (wherein variable groups are, unless otherwise specified, as defined in formula (I)) comprises of~~ comprising:

~~Process 1)~~ a) reacting a compound of formula (II):



(II)


with a compound of formula (III):



(III)

wherein L is a displaceable group;


~~Process 2)~~ b) reacting an acid of formula (IV):



(IV)


or an activated derivative thereof;

with an amine of formula (V):




(V)

~~Process 3):~~ c) reacting an acid of formula (VI):




(VI)

or an activated derivative thereof, with an amine of formula (VII):




(VII)


~~Process 4~~: d) reducing a compound of formula (VIII):



~~Process 5~~: e) reacting a compound of formula (IX):




with a compound of formula (X):



wherein L is a displaceable group;


~~Process 6~~: f) reacting a compound of formula (XI):



(XI)


wherein L is a displaceable group;

with a compound of formula (XII):



(XII)

~~Process 7~~: g) De-esterifying a compound of formula (XIII)



(XIII)

wherein the group C(O)OR is an ester group; and

wherein:

R¹ is hydrogen, C₁₋₆alkyl, C₃₋₆cycloalkyl or aryl; wherein said C₁₋₆alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, carbamoyl, carboxy, C₁₋₆alkoxy, N-(C₁₋₆alkyl)amino, N,N-(C₁₋₆alkyl)₂amino, C₁-C₆ alkylcarbonylamino, C₁₋₆alkylS(O)_a wherein a is 0-2, C₃₋₆cycloalkyl or aryl; and wherein any aryl group may be

optionally substituted by one or two substituents selected from halo, hydroxy, C₁₋₆alkyl or C₁₋₆alkoxy;

R² and R⁵ are independently hydrogen, a branched or unbranched C₁₋₆alkyl, C₃₋₆cycloalkyl or aryl; wherein said C₁₋₆alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, cyano, carbamoyl, carboxy, C₁₋₆alkoxy, aryl C₁₋₆alkoxy, (C₁₋₄)₃Si, N-(C₁₋₆alkyl)amino, N,N-(C₁₋₆alkyl)₂amino, C₁₋₆alkylS(O)_a, C₃₋₆cycloalkyl, aryl or aryl C₁₋₆alkylS(O)_a, wherein a is 0-2; and wherein any aryl group may be optionally substituted by one or two substituents selected from halo, hydroxy, C₁₋₆alkyl or C₁₋₆alkoxy;

R³ is hydrogen, alkyl, halo, C₁₋₆alkoxy or C₁₋₆alkylS-;

R⁴ is hydrogen, C₁₋₆alkyl, halo or C₁₋₆alkoxy;

R⁶ is hydrogen, C₁₋₆alkyl, or arylC₁₋₆alkyl;

wherein R⁵ and R² may form a ring with 2-7 carbon atoms and wherein R⁶ and R² may form a ring with 3-6 carbon atoms; and

L is a displaceable group;

and thereafter ~~if necessary or desirable~~ optionally:

- i) converting a compound of the formula (I) into another compound of the formula (I);
- ii) removing any protecting groups;
- iii) forming a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug; or
- iv) separating two or more enantiomers.

~~L is a displaceable group, suitable values for L are for example, a halogeno or sulphonyloxy group, for example a chloro, bromo, methanesulphonyloxy or toluene 4 sulphonyloxy group.~~

~~C(O)OR is an ester group, suitable values for C(O)OR are methoxycarbonyl, ethoxycarbonyl, *t* butoxycarbonyl and benzyloxycarbonyl.~~

21. (new) A method of treating or preventing a hyperlipidemic condition comprising the administration of an effective amount of a compound according to claim 12 to a mammal in need thereof.

22. (new) A method of treating or preventing atherosclerosis comprising the administration of an effective amount of a compound according to claim 12 to a mammal in need thereof.
23. (new) A method for treating or preventing Alzheimers' disease comprising the administration of an effective amount of a compound according to claim 12 to a mammal in need thereof.
24. (new) A method for treating or preventing a cholesterol associated tumor comprising the administration of an effective amount of a compound according to claim 12 to a mammal in need thereof.
25. (new) A pharmaceutical formulation comprising a compound according to claim 12 in admixture with a pharmaceutically acceptable adjuvant, diluent and/or carrier.
26. (new) A process according to claim 20 wherein L is a halogen or sulphonyloxy group.
27. (new) A process according to claim 26 wherein L is a chloro, bromo, methanesulphonyloxy or toluene-4-sulphonyloxy group.
28. (new) A process according to claim 20 wherein the C(O)OR ester group is methoxycarbonyl, ethoxycarbonyl, *t*-butoxycarbonyl, or benzyloxycarbonyl.